

COMBINATORICS OF LINEAR ITERATED FUNCTION SYSTEMS WITH OVERLAPS

NIKITA SIDOROV

ABSTRACT. Let $p_0, \dots, p_{m-1}$ be points in $\mathbb{R}^d$, and let $\{f_j\}_{j=0}^{m-1}$ be a one-parameter family of similitudes of $\mathbb{R}^d$:

$$f_j(x) = \lambda x + (1 - \lambda)p_j, \quad j = 0, \dots, m-1,$$

where $\lambda \in (0, 1)$ is our parameter. Then, as is well known, there exists a unique self-similar attractor S_λ satisfying $S_\lambda = \bigcup_{j=0}^{m-1} f_j(S_\lambda)$. Each $x \in S_\lambda$ has at least one address $(i_1, i_2, \dots) \in \prod_1^\infty \{0, 1, \dots, m-1\}$, i.e., $\lim_n f_{i_1} f_{i_2} \dots f_{i_n}(0) = x$.

We show that for λ sufficiently close to 1, each $x \in S_\lambda \setminus \{p_0, \dots, p_{m-1}\}$ has $2^{\aleph_0}$ different addresses. If λ is not too close to 1, then we can still have an overlap, but there exist x 's which have a unique address. However, we prove that almost every $x \in S_\lambda$ has $2^{\aleph_0}$ addresses, provided S_λ contains no holes and at least one proper overlap. We apply these results to the case of expansions with deleted digits.

Furthermore, we give sharp sufficient conditions for the Open Set Condition to fail and for the attractor to have no holes.

These results are generalisations of the corresponding one-dimensional results, however most proofs are different.

1. ONE-DIMENSIONAL CASE: AN OVERVIEW

The purpose of this paper is to generalise certain results concerning a one-dimensional model considered in [7, 13, 14] so we first describe this model. Let $\lambda \in (1/2, 1)$ be our parameter. Consider a pair of similitudes of $I = [0, 1]$:

$$\begin{aligned} f_0(x) &= \lambda x, \\ f_1(x) &= \lambda x + 1 - \lambda. \end{aligned}$$

They constitute an *iterated function system* (IFS). More precisely, choose 0 as a starting point, and for any sequence $(\varepsilon_1, \varepsilon_2, \dots)$ of 0s and 1s:

$$x = \lim_{N \rightarrow +\infty} f_{\varepsilon_1} \dots f_{\varepsilon_N}(0),$$

a *forward iteration* (the limit is independent of our choice of the starting point). The set of all x 's that are representable in such a form, is called the *attractor* of an IFS. As is well known, in this case $\lambda > 1/2$ yields the attractor $[0, 1]$.

Date: February 2, 2008.

2000 Mathematics Subject Classification. 60J05.

Key words and phrases. Iterated function system, IFS, address, beta-expansion.

Unlike a general IFS, in this model any composition of f_0 and f_1 can be given in a very simple form:

$$f_{\varepsilon_1} \dots f_{\varepsilon_N}(0) = (\lambda^{-1} - 1) \sum_{n=1}^N \varepsilon_n \lambda^n,$$

whence

$$x = \lim_{N \rightarrow \infty} \sum_{n=1}^N \varepsilon_n \lambda^n = (\lambda^{-1} - 1) \sum_{n=1}^{\infty} \varepsilon_n \lambda^n$$

(sometimes called a λ -expansion of x). Since $\lambda > 1/2$, we have a proper *overlap*

$$f_0(I) \cap f_1(I) = [1 - \lambda, \lambda],$$

so one might expect a typical $x \in (0, 1)$ to have infinitely many distinct λ -expansions (= *addresses*) $(\varepsilon_1, \varepsilon_2, \dots) \in \prod_1^{\infty} \{0, 1\}$ – which indeed proves to be the case.

More precisely, put

$$\mathfrak{R}_\lambda(x) = \left\{ (\varepsilon_n)_1^\infty : x = (\lambda^{-1} - 1) \sum_{n=1}^{\infty} \varepsilon_n \lambda^n \right\}.$$

The following important result regarding $\mathfrak{R}_\lambda(x)$ has been obtained by Erdős, Joó and Komornik:

Theorem 1.1. [5] *If $\lambda > g = \frac{\sqrt{5}-1}{2} = 0.618\dots$, then $\mathfrak{R}_\lambda(x)$ has the cardinality of the continuum for each $x \in (0, 1)$.*

Moreover, the golden ratio g proves to be a sharp constant in the previous theorem. Nonetheless, the following metric result holds for $\lambda \in (1/2, g]$:

Theorem 1.2. [13] *For any $\lambda \in (1/2, g]$ the cardinality of $\mathfrak{R}_\lambda(x)$ is the continuum for Lebesgue-a.e. $x \in (0, 1)$.*

Put

$$\mathcal{U}_\lambda = \left\{ x \in (0, 1) \mid (\varepsilon_n)_1^\infty : x = (\lambda^{-1} - 1) \sum_{n=1}^{\infty} \varepsilon_n \lambda^n \right\}$$

(the *set of uniqueness*). By Theorem 1.1, $\mathcal{U}_\lambda = \emptyset$ if $\lambda > g$.

Theorem 1.3. [7] *The set $\mathcal{U}_\lambda$ is:*

- *countable for $\lambda \in (\lambda_*, g)$;*
- *uncountable of zero Hausdorff dimension if $\lambda = \lambda_*$; and*
- *a set of positive Hausdorff dimension for $\lambda \in (1/2, \lambda_*)$.*

Here $\lambda_* = 0.559525\dots$ denote the (transcendental) Komornik-Loreti constant introduced in [9].

The purpose of this paper is to generalise some of these results to linear IFSs in higher dimensions.

2. MULTIDIMENSIONAL CASE: AN ANALOGUE OF THEOREM 1.1

Let $\mathbf{p}_0, \dots, \mathbf{p}_{m-1}$ be distinct points in $\mathbb{R}^d$, and let $\{f_j\}_{j=0}^{m-1}$ be a one-parameter family of similitudes of $\mathbb{R}^d$:

$$(2.1) \quad f_j(\mathbf{x}) = \lambda \mathbf{x} + (1 - \lambda) \mathbf{p}_j, \quad j = 0, \dots, m-1,$$

where $\lambda \in (0, 1)$ is our parameter¹.

Then, as is well known, there exists a unique self-similar *attractor* S_λ satisfying

$$S_\lambda = \bigcup_{j=0}^{m-1} f_j(S_\lambda).$$

Put $\mathcal{A} = \{0, \dots, m-1\}$. Similarly to the one-dimensional model, every $\mathbf{x} \in S_\lambda$ has at least one *address*, i.e., a sequence $(i_1, i_2, \dots) \in \mathcal{A}^{\mathbb{N}}$ such that

$$\begin{aligned} \mathbf{x} &= \lim_{n \rightarrow +\infty} f_{i_1} \dots f_{i_n}(\mathbf{x}_0) \\ &= (\lambda^{-1} - 1) \sum_{n=1}^{\infty} \lambda^n \mathbf{p}_{i_n}, \end{aligned}$$

where $\mathbf{x}_0 \in \mathbb{R}^d$ is arbitrary. We assume the dimension of the convex hull of $\{\mathbf{p}_0, \dots, \mathbf{p}_{m-1}\}$ to be equal to d . (Otherwise we embed S_λ into $\mathbb{R}^{d'}$ with $d' < d$.)

Recall that an IFS is said to satisfy the Open Set Condition (OSC) if there exists an open set $O \subset \mathbb{R}^d$ such that

$$O = \bigcup_{j=0}^{m-1} f_j(O),$$

with the union being disjoint. Loosely speaking, the OSC means that the images $f_j(\Omega)$ do not intersect properly, where Ω is the convex hull of the $\mathbf{p}_j$. Virtually all famous IFS-generated fractals (the Sierpiński gasket, Sierpiński carpet, von Koch curve, etc.) originate from IFSs that satisfy the OSC.

We will be interested in IFSs which do *not* satisfy the OSC. Here is a simple sufficient condition:

Proposition 2.1. *If $\lambda > m^{-1/d}$, then the OSC is not satisfied.*

Proof. Assume there exists an open set O which satisfies the definition. Since $f_j(O) \subset O$ and the images are disjoint, we have $f_i f_j(O) \cap f_{i'} f_{j'}(O) = \emptyset$ if $(i, j) \neq (i', j')$, and by induction, $f_{i_1} \dots f_{i_n}(O) \cap f_{j_1} \dots f_{j_n}(O) = \emptyset$ provided $(i_1, \dots, i_n) \neq (j_1, \dots, j_n)$.

Since $\mathbb{L}_d(f_{i_1} \dots f_{i_n}(O)) = \lambda^{dn} \mathbb{L}_d(O)$ (where $\mathbb{L}_d$ denotes the d -dimensional Lebesgue measure), and the number of different words of length n is m^n , the pigeonhole principle yields a contradiction with $\lambda > m^{-1/d}$. $\square$

¹To simplify our notation, we have decided to avoid notation like $f_j^{(\lambda)}$, since there is never really any confusion regarding which λ is considered at a given moment.

Example 2.2. Let p_0, p_1, p_2 be three noncollinear points in $\mathbb{R}^2$ – vertices of a triangle Δ . Consider the IFS

$$f_j(x) = \lambda x + (1 - \lambda)p_j, \quad j = 0, 1, 2,$$

and, following [1], we denote the attractor by $\mathcal{S}_\lambda$, i.e.,

$$\mathcal{S}_\lambda = \bigcup_{j=0}^2 f_j(\mathcal{S}_\lambda).$$

Note that for $\lambda = 1/2$ the set $\mathcal{S}_\lambda$ is the famous Sierpiński gasket. If $\lambda \leq 1/2$, then the IFS does satisfy the OSC, and for $\lambda \geq 2/3$ we have $\mathcal{S}_\lambda = \Delta$, i.e., $\mathcal{S}_\lambda$ contains no holes. If $\lambda > 1/2$, then we have a *proper overlap*, i.e., $f_i(\Delta) \cap f_j(\Delta)$ has a nonempty interior.

Remark 2.3. For the triangular case a more delicate argument allows one to show that $\lambda > 1/2$ implies the failure of the OSC (instead of $\lambda > 1/\sqrt{3}$ provided by Proposition 2.1) – see [1, Proposition 3.9].

Return to the general case. Put

$$\Omega = \text{conv}(p_0, \dots, p_{m-1}).$$

Clearly, $\mathcal{S}_\lambda \subset \Omega$. We give a universal sufficient condition for Ω to have no holes.

Proposition 2.4. *If $\lambda \geq d/(d+1)$, then $\mathcal{S}_\lambda = \Omega$, i.e., our attractor has no holes.*

Proof. Let $p_0, \dots, p_{k-1}$ be the vertices of Ω (with $k \leq m$) and let $\mathcal{F}_1, \dots, \mathcal{F}_h$ denote its $(d-1)$ -dimensional faces. Notice that $k \geq d+1$, and if $k = d+1$, we have a simplex for which the claim is proved in [1]. Assume $k > d+1$.

For $x \in \Omega$ we denote its distance to $\mathcal{F}_i$ by x_i . Then adding the volumes of the pyramids with the vertex x and the bases $\mathcal{F}_i$ yields

$$(2.2) \quad \frac{1}{d} \sum_{i=1}^h \mathbb{L}_{d-1}(\mathcal{F}_i) \cdot x_i = \mathbb{L}_d(\Omega).$$

It suffices to show that $\Omega = \bigcup_{j=0}^{k-1} f_j(\Omega)$. Assume, on the contrary, that there exists $x \in \Omega \setminus \bigcup_{j=0}^{k-1} f_j(\Omega)$. Since $f_j(\Omega) = \lambda\Omega + (1-\lambda)p_j$, we have

$$(2.3) \quad \frac{x_i}{\text{dist}(p_j, \mathcal{F}_i)} < 1 - \lambda, \quad i = 1, \dots, h.$$

for all $j \in \{0, \dots, k-1\}$ such that $p_j \notin \mathcal{F}_i$. Put

$$\alpha_i = \max_{0 \leq j \leq k-1} \text{dist}(p_j, \mathcal{F}_i) \quad i = 1, \dots, h.$$

Then by (2.2) and (2.3),

$$\lambda < 1 - \frac{d\mathbb{L}_d(\Omega)}{\sum_{i=1}^h \alpha_i \mathbb{L}_{d-1}(\mathcal{F}_i)}.$$

To complete the proof, it suffices to show that

$$(2.4) \quad \sum_{i=1}^h \alpha_i \mathbb{L}_{d-1}(\mathcal{F}_i) \leq d(d+1) \mathbb{L}_d(\Omega).$$

Consider the family of affine copies of Ω with the ratio $d/(d+1)$, i.e.,

$$\left\{ \frac{d}{d+1} \Omega + \frac{1}{d} \mathbf{p}_j \mid j \in \{0, 1, \dots, k-1\} \right\}.$$

Let $0 \leq j_1 < \dots < j_{d+1} \leq k-1$ and let $\Delta_{j_1 \dots j_{d+1}} = \text{conv}(\mathbf{p}_{j_1}, \dots, \mathbf{p}_{j_{d+1}})$, a d -dimensional simplex. We have

$$\bigcap_{r=1}^{d+1} \left(\frac{d}{d+1} \Delta_{j_1 \dots j_{d+1}} + \frac{1}{d} \mathbf{p}_{j_r} \right) \neq \emptyset$$

(the intersection contains the centre of mass of $\Delta_{j_1 \dots j_{d+1}}$), whence

$$\bigcap_{r=1}^{d+1} \left(\frac{d}{d+1} \Omega + \frac{1}{d} \mathbf{p}_{j_r} \right) \neq \emptyset$$

for any $(d+1)$ -tuple, as $\Delta_{j_1 \dots j_{d+1}} \subset \Omega$. Hence, by Helly's theorem (see, e.g., [4]), there exists

$$\mathbf{z} \in \bigcap_{j=0}^{k-1} \left(\frac{d}{d+1} \Omega + \frac{1}{d} \mathbf{p}_j \right).$$

By our construction, the point $\mathbf{z}$ has the following property: if $\mathbf{y} \in \partial\Omega$, and $\mathbf{z} \in [\mathbf{p}_j, \mathbf{y}]$ for some $j \in \{0, 1, \dots, k-1\}$, then $|\mathbf{z}, \mathbf{y}| \geq \frac{1}{d+1} |\mathbf{p}_j, \mathbf{y}|$. Therefore, $\text{dist}(\mathbf{p}_j, \mathcal{F}_i) \leq (d+1) \cdot \text{dist}(\mathbf{z}, \mathcal{F}_i)$ for all $j \in \{0, \dots, k-1\}$, whence by definition, $\alpha_i \leq (d+1) \cdot \text{dist}(\mathbf{z}, \mathcal{F}_i)$. Consequently,

$$\sum_{i=1}^h \alpha_i \mathbb{L}_{d-1}(\mathcal{F}_i) \leq (d+1) \sum_{i=1}^h \text{dist}(\mathbf{z}, \mathcal{F}_i) \cdot \mathbb{L}_{d-1}(\mathcal{F}_i),$$

and to obtain (2.4), it suffices to note that the volume of Ω equals the sum of the volumes of pyramids whose vertex is $\mathbf{z}$, i.e.,

$$\mathbb{L}_d(\Omega) = \frac{1}{d} \sum_{i=1}^h \text{dist}(\mathbf{z}, \mathcal{F}_i) \cdot \mathbb{L}_{d-1}(\mathcal{F}_i).$$

□

Lemma 2.5. Assume λ is such that $S_\lambda = \Omega$ and suppose $i, j \in \mathcal{A}$ are such that $\Omega_{ij} := f_i(\Omega) \cap f_j(\Omega)$ has a nonempty interior.

Then each $\mathbf{x} \in \Omega_{ij}$ has at least one address beginning with i and at least one address beginning with j .

Proof. Since $\mathbf{x} \in f_i(\Omega)$, there exists $\mathbf{x}' \in \Omega$ such that $f_i(\mathbf{x}') = \mathbf{x}$. By our assumption, $\mathbf{x}' \in S_\lambda$, whence $\mathbf{x}' = \lim_n f_{i_2} \dots f_{i_n}(\mathbf{x}_0)$. Therefore, in view of the continuity of f_i , we have $\mathbf{x} = \lim_n f_i f_{i_2} \dots f_{i_n}(\mathbf{x}_0)$. The same argument applies to j . □

Remark 2.6. The condition $S_\lambda = \Omega$, generally speaking, cannot be dropped; for instance, one can show that in the triangular case, if $\lambda \in (0.65, 2/3)$, then we have a hole whose image under one of the maps lies in S_λ – see [1, Proposition 3.7].

Theorem 2.7. *For each $p_0, \dots, p_{m-1}$ there exists $\lambda_0 < 1$ such that for any $\lambda \in (\lambda_0, 1)$,*

- (1) *There are no holes, i.e., $S_\lambda = \Omega$;*
- (2) *Each point $x \in \Omega$, except when x is a vertex of Ω , has 2^{N_0} distinct addresses.*

Proof. Assume that λ is large enough to ensure $S_\lambda = \Omega$. We will show that each x under consideration has a continuum of addresses $(i_1, i_2, \dots) \in \mathcal{A}'$, where $\mathcal{A}'$ is the set of indices i such that $p_i \in \partial\Omega$.

The idea of the proof is to use the multivalued *inverse map* $T_\lambda = \{f_0, \dots, f_{m-1}\}^{-1}$. More precisely, put

$$\Omega_i = f_i(\Omega) \setminus \bigcup_{j \neq i} f_j(\Omega).$$

Clearly, if $\Omega_i \ni x \sim (i_1, i_2, \dots)$, then necessarily $i_1 = i$. Conversely, if a point $x \notin \Omega_i$ for any i , then by Lemma 2.5, there is a choice for the first symbol of its address.

Let $x \in \Omega_i \setminus \{p_i\}$; note that shifting its address $(i, i_2, i_3, \dots)$ yields $(i_2, i_3, \dots)$, which in Ω corresponds to applying f_i^{-1} . Let λ_0 be such that $f_i^{-1}(\Omega_i) \cap \Omega_j = \emptyset$ for all $i \neq j$ and all $\lambda > \lambda_0$.

Thus, by f_i^{-1} being expanding on $\Omega_i \setminus \{p_i\}$, we conclude that there exists $k \geq 1$ such that

$$f_i^{-k}(x) \in \Omega_i,$$

$$x' = f_i^{-k-1}(x) \in \Omega \setminus \bigcup_{i=0}^{m-1} \Omega_i.$$

By our construction, x' has at least two addresses; consider its two shifts, x'' and x''' , say. Either $x'' \in \Omega \setminus \bigcup_{i=0}^{m-1} \Omega_i$ and thus, has at least two addresses itself or it belongs to Ω_j for some j (and obviously, is not equal to p_j) and, similarly to the above, we shift its address until it falls into $\Omega \setminus \bigcup_{i=0}^{m-1} \Omega_i$. Hence any $x \in \Omega$ that is not one of its vertices, has 2^{N_0} distinct addresses. $\square$

Remark 2.8. Note that for the triangular case the sharp constant is $\lambda_0 \approx 0.68233$, the unique positive root of $x + x^3 = 1$ – see Theorem 4.1.

3. MULTIDIMENSIONAL CASE: GENERIC BEHAVIOUR

3.1. General theory: Lebesgue measure. Let U_λ denote the set of $x \in S_\lambda$ having a unique address, and $R_\lambda(x)$ denote the set of all addresses of a given $x \in S_\lambda$.

Lemma 3.1. *Put*

$$V_\lambda = \{x \in S_\lambda : \text{card } R_\lambda(x) < 2^{N_0}\}.$$

Then $\dim_H V_\lambda = \dim_H U_\lambda$ (where $\dim_H$ denotes Hausdorff dimension).

Proof. We are going to exploit the idea of *branching* introduced in [14]. Let $x \in S_\lambda$ have at least two addresses; then there exists the smallest $n \geq 0$ such that $x \sim (i_1, \dots, i_n, i_{n+1}, \dots)$




FIGURE 1. Branching and bifurcations.

and $\mathbf{x} \sim (i_1, \dots, i_n, i'_{n+1}, \dots)$ with $i_{n+1} \neq i'_{n+1}$. We may depict this *bifurcation* as is shown in Fig. 1.

Assume that $\mathbf{x}$ has less than a continuum of distinct addresses. Then, inevitably, one of the branches at some point ceases to bifurcate. In other words, there exists $(i_n)_1^\infty \in \mathcal{A}^\mathbb{N}$ and $N \in \mathbb{N}$ such that $\mathbf{x} \sim (i_1, i_2, \dots)$ and $\mathbf{x}' \sim (i_N, i_{N+1}, \dots) \in U_\lambda$. Hence

$$(3.1) \quad V_\lambda \subset \bigcup_{(i_1, \dots, i_N) \in \mathcal{A}^N} f_{i_1} \dots f_{i_N}(U_\lambda).$$

Since the f_i are linear, (3.1) implies $\dim_H V_\lambda \leq \dim_H U_\lambda$, and the inverse inequality is trivial. $\square$

Remark 3.2. Note that if λ_0 in Theorem 2.7 is sharp, we always have a nonempty set of uniqueness for $\lambda < \lambda_0$, because, as we know from the branching argument, the existence of $\mathbf{x}$ with less than a continuum of addresses implies the existence of $\mathbf{x}'$ with a unique address. For an example see Section 4.

Our goal is to show that, similarly to the one-dimensional case, if there are no holes and at least one proper overlap, then a.e. $\mathbf{x}$ has a continuum of addresses. We need an auxiliary claim from dimension theory:

Lemma 3.3. *Let $A \subset \mathbb{R}^d$ be such that there exists a positive constant $\delta > 0$ such that for an arbitrary cube $C \subset \mathbb{R}^d$ which intersects A , one can find a cube $C_0 \subset C$ such that $L_d(C_0) \geq \delta L_d(C)$ and $C_0 \cap A = \emptyset$.*

Then $\dim_H A < d$.

Proof. Since $\dim_H A \leq \underline{\dim}_B A$ (lower box-counting dimension), it suffices to show that $\underline{\dim}_B A < d$. Recall that there are various definitions of $\underline{\dim}_B$ (see [6, Chapter 3.1]) and in particular, the one which involves mesh cubes which we will use.

More precisely, a cube of the form $[m_1\varepsilon, (m_1 + 1)\varepsilon] \times \cdots \times [m_d\varepsilon, (m_d + 1)\varepsilon]$ for some $\varepsilon > 0$ and $m_1, \dots, m_d$ integers, is called an ε -mesh cube. Let $\mathcal{N}_\varepsilon(A)$ denote the number of ε -mesh cubes which intersect A . Then

$$\underline{\dim}_B A = \liminf_{\varepsilon \rightarrow 0} \frac{\log \mathcal{N}_\varepsilon(A)}{\log(1/\varepsilon)}.$$

It is obvious that our condition implies that there exists $M \in \mathbb{N}$ such that for any ε -mesh cube C which intersects A , there exists an ε/M -mesh cube $C_0 \subset C$ which doesn't. Hence

$$\mathcal{N}_{\varepsilon/M}(A) \leq (M^d - 1)\mathcal{N}_\varepsilon(A),$$

which implies

$$\mathcal{N}_{M^{-n}}(A) \leq (M^d - 1)^n \cdot \mathcal{N}_1(A).$$

Consequently,

$$\liminf_{\varepsilon \rightarrow 0} \frac{\log \mathcal{N}_\varepsilon(A)}{\log 1/\varepsilon} \leq \frac{\log(M^d - 1)}{\log M} < d.$$

□

Now we are ready to prove a key technical lemma.

Lemma 3.4. *Let $\{F_j\}_{j=0}^{L-1}$ be a finite family of linear contractions of $\mathbb{R}^d$ with the same contraction ratio $\beta \in (0, 1)$. Let $\Omega \subset \mathbb{R}^d$ be a polyhedron of dimension d , and assume $F_j(\Omega) \subset \Omega$ for all $j = 0, \dots, L-1$, and $\Omega = \bigcup_{j=0}^{L-1} F_j(\Omega)$ (no holes).*

Put

$$(3.2) \quad W_n = \bigcup_{\substack{(j_1, \dots, j_n) \in \{0, \dots, L-1\}^n: \\ \exists k \in \{1, \dots, n\}: j_k = 0}} F_{j_1} \dots F_{j_n}(\Omega).$$

Then $W_n \subset W_{n+1}$ for all $n \geq 1$, and $\mathcal{L}_d(\Omega \setminus W) = 0$, where $W = \bigcup_{n \geq 1} W_n$. Furthermore, $\dim_H(\Omega \setminus W) < d$.

Proof. Our first goal is to show that $W_n \subset W_{n+1}$ for all $n \geq 1$. Since $\Omega = \bigcup_{j=0}^{L-1} F_j(\Omega)$, we have by induction,

$$\bigcup_{(j_1, \dots, j_k) \in \{0, \dots, L-1\}^k} F_{j_1} \dots F_{j_k}(\Omega) = \Omega,$$

whence for any $k < n$,

$$\bigcup_{(j_1, \dots, j_n) \in \{0, \dots, L-1\}^n} F_{j_1} \dots F_{j_k} F_0 F_{j_{k+1}} \dots F_{j_n}(\Omega) = \bigcup_{(j_1, \dots, j_k) \in \{0, \dots, L-1\}^k} F_{j_1} \dots F_{j_k} F_0(\Omega).$$

Hence by definition, $W_1 = F_0(\Omega)$ and

$$W_n = \left(\bigcup_{k=1}^{n-1} \bigcup_{(j_1, \dots, j_k) \in \{0, \dots, L-1\}^k} F_{j_1} \dots F_{j_k} F_0(\Omega) \right) \cup F_0(\Omega), \quad n \geq 2.$$

Consequently,

$$W_{n+1} = W_n \cup \bigcup_{(j_1, \dots, j_n) \in \{0, \dots, L-1\}^n} F_{j_1} \dots F_{j_n} F_0(\Omega), \quad n \geq 1,$$

whence $W_n \subset W_{n+1}$.

By Lemma 3.3, to show that $\dim_H(\Omega \setminus W) < d$, it suffices to demonstrate that there exists a positive constant $\delta = \delta(\Omega, \beta) > 0$ such that given an arbitrary cube $\mathcal{C} \subset \Omega$, one can find a cube $\mathcal{C}_0 \subset \mathcal{C}$ such that $\mathbb{L}_d(\mathcal{C}_0) \geq \delta \mathbb{L}_d(\mathcal{C})$ and $\mathcal{C}_0 \cap (\Omega \setminus W) = \emptyset$.

So we choose an arbitrary cube $\mathcal{C} \subset \Omega$ and denote the length of its edge by κ . Let ξ denote the centre of $\mathcal{C}$; since Ω has no holes, $\xi = \lim_{r \rightarrow \infty} F_{j_1} \dots F_{j_r}(\Omega)$ (the limit in the Hausdorff metric). Hence there exists a unique N such that $F_{j_1} \dots F_{j_{N-1}}(\Omega) \not\subset \mathcal{C}$, and $F_{j_1} \dots F_{j_N}(\Omega) \subset \mathcal{C}$.

Notice that since $\xi \in F_{j_1} \dots F_{j_{N-1}}(\Omega)$ and $F_{j_1} \dots F_{j_{N-1}}(\Omega) \not\subset \mathcal{C}$, we have

$$\text{diam } F_{j_1} \dots F_{j_{N-1}}(\Omega) \geq \text{dist}(\xi, \partial \mathcal{C}) = \kappa/2.$$

Put

$$\nu = \frac{\text{diam}(\Omega)}{(\mathbb{L}_d(\Omega))^{1/d}}.$$

Then

$$\begin{aligned} \mathbb{L}_d(\mathcal{C})^{1/d} &= \kappa \leq 2 \text{diam } F_{j_1} \dots F_{j_{N-1}}(\Omega) = 2\beta^{N-1} \text{diam}(\Omega) \\ &= \frac{2\nu}{\beta} \cdot \beta^N (\mathbb{L}_d(\Omega))^{1/d} = \frac{2\nu}{\beta} \cdot (\mathbb{L}_d(F_{j_1} \dots F_{j_{N-1}}(\Omega)))^{1/d}, \end{aligned}$$

whence

$$(3.3) \quad \frac{\mathbb{L}_d(F_{j_1} \dots F_{j_N}(\Omega))}{\mathbb{L}_d(\mathcal{C})} \geq c > 0,$$

where $c = (\beta/2\nu)^d$, i.e., c depends only on the shape of Ω and on the contraction ratio, but not on $\mathcal{C}$ itself.

Put $\Omega_0 = F_{j_1} \dots F_{j_N} F_0(\Omega)$; by (3.3), $\mathbb{L}_d(\Omega_0) \geq \beta c \cdot \mathbb{L}_d(\mathcal{C})$. We can find a cube $\mathcal{C}_1 \subset \Omega$ such that the ratio of their volumes equals $\gamma > 0$. Since Ω_0 is similar to Ω , we put $\mathcal{C}_0 = F_{j_1} \dots F_{j_N} F_0(\mathcal{C}_1) \subset \Omega_0$ and obtain

$$\mathbb{L}_d(\mathcal{C}_0) \geq \beta \gamma c \cdot \mathbb{L}_d(\mathcal{C}),$$

where $\delta := \beta \gamma c$ is independent of $\mathcal{C}$. Furthermore, Ω_0 (and consequently, $\mathcal{C}_0$) has an empty intersection with $\Omega \setminus W_{N+1}$, whence $\mathcal{C}_0 \cap (\Omega \setminus W) = \emptyset$ as well, and we are done. $\square$

Theorem 3.5. Assume

- $S_\lambda = \Omega$, i.e., there are no holes;
- there exist $i, k \in \mathcal{A}$ such that a vertex of $f_k(\Omega)$ belongs to the interior of $f_i(\Omega)$.

Then $\mathbb{L}_d$ -a.e. $x \in \Omega$ has $2^{\aleph_0}$ distinct addresses, and the exceptional set V_λ has Hausdorff dimension strictly less than d .

Remark 3.6. If $d \leq 2$, it suffices to assume that $f_i(\Omega) \cap f_k(\Omega)$ has a nonempty interior, since if two convex polygons (or intervals) intersect properly, then it is obvious that there exists a vertex of one which lies in the interior of the other. For $d \geq 3$ this is not always the case.

Proof. By our assumption, there exists $j \in \mathcal{A}$ such that $f_k(\mathbf{p}_j) \in \text{int}(f_i(\Omega))$. Hence there exists $\ell \in \mathbb{N}$ such that $f_i f_j^{\ell-1}(\Omega) \subset f_i(\Omega) \cap f_k(\Omega)$. Since $S_\lambda = \Omega$, Lemma 2.5 implies that any $x \in f_i f_j^{\ell-1}(\Omega)$ has at least two different addresses.

Put $L = m^\ell$ and define $\{F_0, \dots, F_{L-1}\} = \{f_{i_1} \dots f_{i_\ell} \mid (i_1, \dots, i_\ell) \in \mathcal{A}^\ell\}$ with $F_0 = f_i f_j^{\ell-1}$. By the above, each $x \in F_{j_1} \dots F_{j_{k-1}} F_0 F_{j_{k+1}} \dots F_n(\Omega)$ has at least two different addresses, whence $U_\lambda \subset \Omega \setminus W$, where $W = \bigcup_n W_n$ and W_n is given by (3.2).

Hence by Lemma 3.4, $\dim_H(U_\lambda) < d$, whence by Lemma 3.1, $\dim_H(V_\lambda) < d$, which is the claim of the theorem. $\square$

Remark 3.7. If λ is sufficiently close to the critical value λ_0 (see the previous section), then one could expect the exceptional set V_λ to be countable (similarly to the one-dimensional case).

3.2. Application: λ -expansions with deleted digits. Expansions of real numbers in non-integer bases with deleted digits have been studied since the mid-1990s – see, e.g., [8, 12]. The model is as follows: assume $d = 1$ and let $A = \{a_1, \dots, a_m\} \subset \mathbb{R}$ be a “digit” set with $a_1 < \dots < a_m$. Let $x \in \mathbb{R}$ have an expansion of the form

$$(3.4) \quad x = \sum_{n=1}^{\infty} \varepsilon_n \lambda^n, \quad \varepsilon_n \in A, \quad n \geq 1.$$

It is obvious that $\lambda a_1 / (1 - \lambda) \leq x \leq \lambda a_m / (1 - \lambda)$. M. Pedicini [11] has shown that if

$$(3.5) \quad \max_{1 \leq j \leq m-1} (a_{j+1} - a_j) < \frac{\lambda(a_m - a_1)}{1 - \lambda},$$

then each $x \in [\lambda a_1 / (1 - \lambda), \lambda a_m / (1 - \lambda)]$ has at least one expansion of the form (3.4). Note also that in the recent paper [2] the theory of random and greedy beta-expansions with deleted digits (under the assumption (3.5)) has been developed.

We apply our results from this and the previous section to obtain

Proposition 3.8. (1) *There exists $\lambda_0 = \lambda_0(a_1, \dots, a_m) < 1$ such that for each $\lambda \in (\lambda_0, 1)$ any $x \in (\lambda a_1 / (1 - \lambda), \lambda a_m / (1 - \lambda))$ has $2^{\aleph_0}$ expansions of the form (3.4).*
 (2) *If the condition (3.5) is satisfied, Lebesgue-a.e. $x \in (\lambda a_1 / (1 - \lambda), \lambda a_m / (1 - \lambda))$ has $2^{\aleph_0}$ expansions of the form (3.4), and the exceptional set has Hausdorff dimension strictly less than 1.*

Proof. Put

$$(3.6) \quad f_j(x) = \lambda(x + a_j), \quad 1 \leq j \leq m.$$

Then, as in the standard one-dimensional case (where $a_1 = 0, a_2 = 1$), we have by induction,

$$f_{\varepsilon_1} \dots f_{\varepsilon_n}(x_0) = \lambda^n x_0 + \sum_{k=1}^n \varepsilon_k \lambda^k,$$

whence

$$\lim_{n \rightarrow \infty} f_{\varepsilon_1} \dots f_{\varepsilon_n}(x_0) = \sum_{n=1}^{\infty} \varepsilon_n \lambda^n$$

for any $x_0 \in \mathbb{R}$. Therefore, x has an expansion of the form (3.4) if and only if $x \in S_\lambda$ for the IFS (3.6).

The condition (3.5) ensures that $S_\lambda = \Omega = [\lambda a_1/(1-\lambda), \lambda a_m/(1-\lambda)]$. To prove the first part of the proposition, notice that (3.5) holds for all λ sufficiently close to 1 so Theorem 2.7 is applicable to any x which lies in the interior of Ω . (As $\partial\Omega = \{a_1/(1-\lambda), a_m/(1-\lambda)\}$.)

To prove the second part, we notice that by (3.5),

$$|f_j(\Omega)| = \frac{\lambda^2(a_m - a_1)}{1 - \lambda} > \lambda(a_{j+1} - a_j), \quad 1 \leq j \leq m-1.$$

Hence

$$\begin{aligned} \sum_{j=1}^m |f_j(\Omega)| &= \sum_{j=1}^{m-1} |f_j(\Omega)| + |f_m(\Omega)| \\ &> \lambda(a_m - a_1) + \frac{\lambda^2(a_m - a_1)}{1 - \lambda} \\ &= \frac{\lambda(a_m - a_1)}{1 - \lambda} = |\Omega|, \end{aligned}$$

whence there exists $j \in \{1, \dots, m-1\}$ such that $f_j(\Omega) \cap f_{j+1}(\Omega)$ has a nonempty interior. Thus, we can apply Theorem 3.5 to this setting. $\square$

Remark 3.9. In her PhD dissertation, Anna-Chiara Lai [10] has proved a weaker version of the second claim of Proposition 3.8.

Finally, we prove

Lemma 3.10. *Provided (3.5) is satisfied, the Open Set Condition for the IFS (3.6) fails.*

Proof. We have

$$a_m - a_1 = \sum_{j=1}^{m-1} (a_{j+1} - a_j) < \frac{\lambda(m-1)(a_m - a_1)}{1 - \lambda},$$

whence $\lambda > 1/m$, and we apply Proposition 2.1. $\square$

3.3. General theory: natural measure. In the end of this section we would like to obtain a result similar to Theorem 3.5 for a “natural” measure on S_λ . Let $(p_0, p_1, \dots, p_{m-1})$ be a probability vector with $p_j > 0$ for all j . The *probabilistic IFS* given by the f_i and the p_i is defined as follows: put $\Sigma = \mathcal{A}^{\mathbb{N}}$ and define ρ as the product measure on Σ with equal multipliers $(p_0, p_1, \dots, p_{m-1})$. Let the projection map $\pi : \Sigma \rightarrow \mathbb{R}^d$ be given by the formula

$$\pi(i_1, i_2, \dots) := \lim_{n \rightarrow +\infty} f_{i_1} f_{i_2} \dots f_{i_n}(\mathbf{0}).$$

We define the measure μ on S_λ as the push down measure $\pi(\rho)$. As is well known, $\text{supp}(\mu) = S_\lambda$.

Proposition 3.11. *Under the assumptions of Theorem 3.5, μ -a.e. $x \in \Omega$ has $2^{\aleph_0}$ distinct addresses.*

Proof. Put $\mathbf{w} = ij^{\ell-1}$. By the Birkhoff ergodic theorem applied to the one-sided Bernoulli shift on the measure space (Σ, ρ) ,

$$\rho\{(i_1, i_2, \dots) \in \Sigma \mid \exists k : (i_k, \dots, i_{k+\ell-1}) = \mathbf{w}\} = 1,$$

whence $\mu(U_\lambda) = 0$, because $\mathbf{x} \in U_\lambda$ cannot have an address containing $\mathbf{w}$. All that is left is to show that $\mu(V_\lambda) = 0$ as well.

In view of (3.1), it suffices to show that

$$(3.7) \quad \mu(f_{i_1} \dots f_{i_n}(U_\lambda)) = 0, \quad \forall (i_1, \dots, i_n) \in \mathcal{A}^n.$$

Note that, as is well known (see, e.g., [3]), the self-similarity of the measure μ implies

$$\mu(E) = \sum_{s=0}^{m-1} p_s \cdot \mu(f_s(E)),$$

for any Borel set E . Hence by induction,

$$\mu(E) = \sum_{(i_1, \dots, i_n) \in \mathcal{A}^n} p_{i_1} \dots p_{i_n} \cdot \mu(f_{i_1} \dots f_{i_n}(E)),$$

which implies (3.7). □

Remark 3.12. In fact, to apply the ergodic theorem to the shift on Σ , all we need from ρ is $p_i > 0$ and $p_j > 0$; the other components of the probability vector may equal zero.

The main problem for the future study is to check whether in some cases of IFSs with holes an analogue of Theorem 3.5 still holds. We plan to study this question in our subsequent papers.

4. MAIN EXAMPLE: TRIANGLE

For the triangular case we give an explicit analogue of one-dimensional results mentioned in Section 1. Following [1], we denote the set of uniqueness by $\mathcal{U}_\lambda$.

Theorem 4.1. *Let $\lambda_0 \approx 0.68233$ be the unique positive root of $x^3 + x = 1$. Then*

- (1) *for $\lambda < \lambda_0$, then the set of uniqueness $\mathcal{U}_\lambda$ is nonempty;*
- (2) *if $\lambda \in (\lambda_0, 1)$, then each $x \in \mathcal{S}_\lambda \setminus \{\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2\}$ has $2^{\aleph_0}$ different addresses.*

Proof. (1) Firstly, we introduce a convenient coordinate system for this case suggested in [1]. Without loss of generality, we may assume our triangle Δ to be equilateral. We now identify each point $\mathbf{x} \in \Delta$ with a triple (x, y, z) , where

$$x = \text{dist}(\mathbf{x}, [\mathbf{p}_1, \mathbf{p}_2]), \quad y = \text{dist}(\mathbf{x}, [\mathbf{p}_0, \mathbf{p}_2]), \quad z = \text{dist}(\mathbf{x}, [\mathbf{p}_0, \mathbf{p}_1]),$$

where $[\mathbf{p}_i, \mathbf{p}_j]$ is the edge containing $\mathbf{p}_i$ and $\mathbf{p}_j$. As is well known, $x + y + z$ equals the tripled radius of the inscribed circle, and we choose it to be equal to 1. These coordinates are called *barycentric*. Henceforward we write each $\mathbf{x} \in \mathcal{S}_\lambda$ in barycentric coordinates.

It is shown in [1] that $(x, y, z) \in \mathcal{S}_\lambda$ if and only if there exist three 0-1 sequences $(a_n)_0^\infty$, $(b_n)_0^\infty$ and $(c_n)_0^\infty$ such that

$$\begin{aligned} x &= (1 - \lambda) \sum_{n=0}^{\infty} a_n \lambda^n, \\ y &= (1 - \lambda) \sum_{n=0}^{\infty} b_n \lambda^n, \\ z &= (1 - \lambda) \sum_{n=0}^{\infty} c_n \lambda^n, \end{aligned}$$

with $a_n + b_n + c_n = 1$ for all $n \geq 0$.

We claim that the point

$$(4.1) \quad \pi(\lambda) = \left(\frac{\lambda^2}{1 + \lambda + \lambda^2}, \frac{\lambda}{1 + \lambda + \lambda^2}, \frac{1}{1 + \lambda + \lambda^2} \right)$$

belongs to $\mathcal{U}_\lambda$ provided $\lambda < \lambda_0$. To prove this, it suffices to demonstrate that the system of equations

$$(4.2) \quad \begin{aligned} a_0 + a_1 \lambda + a_2 \lambda^2 + \dots &= \frac{\lambda^2}{1 - \lambda^3}, \\ b_0 + b_1 \lambda + b_2 \lambda^2 + \dots &= \frac{\lambda}{1 - \lambda^3}, \\ c_0 + c_1 \lambda + c_2 \lambda^2 + \dots &= \frac{1}{1 - \lambda^3} \end{aligned}$$

has a unique solution $(a_n)_0^\infty = (001001\dots)$, $(b_n)_0^\infty = (010010\dots)$, $(c_n)_0^\infty = (100100\dots)$. Note first that a_0 cannot be equal to 1 nor can b_0 , because $\lambda < \lambda_0$ implies $1 > \frac{\lambda}{1 - \lambda^3} > \frac{\lambda^2}{1 - \lambda^3}$. Hence $a_0 = b_0 = 0, c_0 = 1$. Similarly, $a_1 = 0$ for the same reason as b_0 , and $c_1 = 0$ as well, because $\lambda > \frac{\lambda^3}{1 - \lambda^3}$. Thus, $b_1 = 1$. Finally, c_2 and b_2 must be equal to 0, whilst $a_2 = 1$.


Thus, we have

$$\begin{aligned} a_3 + a_4 \lambda + a_5 \lambda^2 + \dots &= \frac{\lambda^2}{1 - \lambda^3}, \\ b_3 + b_4 \lambda + b_5 \lambda^2 + \dots &= \frac{\lambda}{1 - \lambda^3}, \\ c_3 + c_4 \lambda + c_5 \lambda^2 + \dots &= \frac{1}{1 - \lambda^3}, \end{aligned}$$

and we can continue the process *ad infinitum*. Therefore, each a_k, b_k and c_k is uniquely determined from the system of equations (4.2), whence $\pi(\lambda) \in \mathcal{U}_\lambda$.

(2) Suppose $\lambda > \lambda_0$. Following the argument of the proof of Theorem 2.7, we introduce the sets

$$\Delta_i := \Delta \setminus \bigcup_{j \neq i} f_j(\Delta)$$

FIGURE 2. Triangular case: $f_0^{-1}(\Gamma_0) \cap (\bigcup_i \Delta_i) = \emptyset$.

(three rhombi). Thus, if $\mathbf{x} \in \mathcal{U}_\lambda$, then necessarily $\mathbf{x} \in \bigcup_i \Delta_i$. Fix $i \in \{0, 1, 2\}$; again, since the shift map on Δ_i , i.e., f_i^{-1} , is expanding on $\Delta_i \setminus \{p_i\}$, eventually $f_i^{-n}(\mathbf{x}) \in \Delta_i$ and $f_i^{-n-1}(\mathbf{x}) \notin \Delta_i$ for some $n \geq 0$, for any $\mathbf{x} \in \Delta_i \setminus \{p_i\}$.

If $f_i^{-n-1}(\mathbf{x}) \notin \bigcup_{j \neq i} \Delta_j$, then $\mathbf{x} \notin \mathcal{U}_\lambda$. Put

$$\Gamma_i = f_i^{-1}(\Delta_j) \cap \Delta_i, \quad j \neq i.$$

In view of the symmetry, the choice of $j \neq i$ is unimportant – see Fig 2. Thus, $\mathcal{U}_\lambda \neq \emptyset$ implies $\bigcup_i \Gamma_i \neq \emptyset$.

Note that the Γ_i are equal for $i = 0, 1, 2$, whence $\bigcup_i \Gamma_i \neq \emptyset \Leftrightarrow \Gamma_0 \neq \emptyset$. The latter is in fact equivalent to $\lambda < 1/\sqrt{2}$. Indeed, we have in barycentric coordinates,

$$\Gamma_0 = \{x < (1 - \lambda)/\lambda, y < 1 - \lambda, z < 1 - \lambda\}.$$

An open triangle $\{x < a, y < b, z < c\}$ is nondegenerate if and only if $a + b + c > 1$. Hence $(1 - \lambda)/\lambda + 2(1 - \lambda) > 1$, which is equivalent to $\lambda < 1/\sqrt{2}$. Thus, $\lambda \in (\lambda_0, 1/\sqrt{2})$.

Assume $\mathbf{x} \in \Gamma_0 \cap \mathcal{U}_\lambda$; then $f_0^{-1}(\mathbf{x})$ has to intersect $\bigcup_i \Delta_i$. It is easy to check that $f_0^{-1}(\Gamma_0) \cap \Delta_0 = \emptyset$, whence, in view of the symmetry, $f_0^{-1}(\Gamma_0) \cap \Delta_1 \neq \emptyset$. We have

$$f_0^{-1}(\Gamma_0) = \left\{ x < \left(\frac{1 - \lambda}{\lambda} \right)^2, y < \frac{1 - \lambda}{\lambda}, z < \frac{1 - \lambda}{\lambda} \right\},$$

$$\Delta_1 = \{x < 1 - \lambda, y < 1 - \lambda\},$$

and we claim that actually, $f_0^{-1}(\Gamma_0)$ lies strictly on the right of Δ_1 – see Fig 2.

Indeed, the coordinates of the point $\mathcal{M}$, the top left corner of the triangle $f_0^{-1}(\Gamma_0)$, are as follows: $\mathcal{M} \left(\left(\frac{1-\lambda}{\lambda} \right)^2, \frac{1-\lambda}{\lambda}, \frac{-1+\lambda+\lambda^2}{\lambda^2} \right)$, and we observe that the inequality $\frac{-1+\lambda+\lambda^2}{\lambda^2} > 1 - \lambda$ is equivalent to $\lambda > \lambda_0$.

Thus, each $x \in \Delta \setminus \{p_0, p_1, p_2\}$ has at least two addresses of the form $x \sim (i_1, \dots, i_n, i_{n+1}, \dots)$ and $x \sim (i_1, \dots, i_n, j_{n+1}, \dots)$ with $j_{n+1} \neq i_{n+1}$ and a “compulsory” prefix $(i_1, \dots, i_n)$, which may be empty (see Fig. 1). Hence x has $2^{\aleph_0}$ different addresses. $\square$

Remark 4.2. One can easily obtain from the proof of the previous theorem that for $\mathcal{U}_{\lambda_0} = \emptyset$ as well, but in fact, the point $\pi(\lambda_0) = (\lambda_0^4, \lambda_0^3, \lambda_0^2)$ has only $\aleph_0$ different addresses. Thus, λ_0 is indeed the full analogue of the golden ratio for the triangular model. We leave the details as an exercise for the reader.

Remark 4.3. Note that if, like in the proof of Theorem 2.7, T_λ denotes the inverse of $\{f_0, f_1, f_2\}$ (well defined on $\bigcup_i \Delta_i$), then $\pi(\lambda)$ given by (4.1) is a period 3 point for T_λ , i.e., $T_\lambda^3 \pi = \pi$. Another 3-cycle is generated by $\pi'(\lambda) = \left(\frac{1}{1+\lambda+\lambda^2}, \frac{\lambda}{1+\lambda+\lambda^2}, \frac{\lambda^2}{1+\lambda+\lambda^2} \right)$, and we conjecture that if $\frac{2}{3} \leq \lambda < \lambda_0$, then $\bigcup_{i=0}^2 \Gamma_i \cap \mathcal{U}_\lambda$ consists of just these 6 points. This would imply that $\mathcal{U}_\lambda$ is countable for this range of parameters.

A full “triangular” analogue of Theorem 1.3 is yet to be determined. In particular, what is the analogue of the Komornik-Loreti constant for the triangular case?

Note that for $\lambda = g = (\sqrt{5} - 1)/2$ the set $\mathcal{U}_\lambda$ is a continuum naturally isomorphic to the space of one-sided 0-1 sequences, and its Hausdorff dimension is $-\log 2 / \log g$ – see [1, Theorem 6.4] and Fig. 5 therein.

Acknowledgment. The author is indebted to F. Petrov for his generous help with the proof of Proposition 2.4 and especially to the anonymous referee for many useful remarks and suggestions.

REFERENCES

- [1] D. Broomhead, J. Montaldi and N. Sidorov, *Golden gaskets: variations on the Sierpinski sieve*, Nonlinearity **17** (2004), 1455–1480.
- [2] K. Dajani and C. Kalle, *Random beta-expansions with deleted digits*, Discr. Cont. Dynam. Systems **18** (2007), 199–217.
- [3] P. Diaconis and D. Freedman, *Iterated random functions*, SIAM Review **41** (1999), 45–76.
- [4] J. Eckhoff, *Helly, Radon, and Caratheodory type theorems*, *Handbook of convex geometry*, Vol. A, B, 389–448, North-Holland, Amsterdam, 1993.
- [5] P. Erdős, I. Joó and V. Komornik, *Characterization of the unique expansions $1 = \sum_{i=1}^{\infty} q^{-n_i}$ and related problems*, Bull. Soc. Math. Fr. **118** (1990), 377–390.
- [6] K. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, John Wiley & Sons, 2003.
- [7] P. Glendinning and N. Sidorov, *Unique representations of real numbers in non-integer bases*, Math. Res. Lett. **8** (2001), 535–543.
- [8] M. Keane, M. Smorodinsky and B. Solomyak, *On the morphology of γ -expansions with deleted digits*, Trans. Amer. Math. Soc. **347** (1995), 955–966.
- [9] V. Komornik and P. Loreti, *Unique developments in non-integer bases*, Amer. Math. Monthly **105** (1998), 636–639.

- [10] A.-Ch. Lai, private communication.
- [11] M. Pedicini, *Greedy expansions and sets with deleted digits*, Theoret. Comp. Sci., **332** (2005), 313–336.
- [12] M. Pollicott and K. Simon, *The Hausdorff dimension of λ -expansions with deleted digits*, Trans. Amer. Math. Soc. **347** (1995), 967–983.
- [13] N. Sidorov, *Almost every number has a continuum of β -expansions*, Amer. Math. Monthly **110** (2003), 838–842.
- [14] N. Sidorov, *Universal β -expansions*, Period. Math. Hungar. **47** (2003), 221–231.

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF MANCHESTER, P.O. BOX 88, SACKVILLE STREET, MANCHESTER M60 1QD, UNITED KINGDOM. E-MAIL: SIDOROV@MANCHESTER.AC.UK